



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07F 9/38	A1	(11) International Publication Number: WO 00/01707 (43) International Publication Date: 13 January 2000 (13.01.00)
<p>(21) International Application Number: PCT/EP99/04587</p> <p>(22) International Filing Date: 2 July 1999 (02.07.99)</p> <p>(30) Priority Data: 1436/98 6 July 1998 (06.07.98) CH</p> <p>(71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).</p> <p>(71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT M.B.H. [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): SIEBENHAAR, Bernd [DE/DE]; Maugenharderstrasse 7, D-79400 Kandern (DE). STUDER, Martin [CH/CH]; Im Zimmerhof 2, CH-4054 Basel (CH).</p> <p>(74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).</p>	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
(54) Title: PREPARATION OF N-PHOSPHONOMETHYLGLYCINE SALTS		
(57) Abstract		
<p>Process for the preparation of N-phosphonomethylglycine salts by reacting a salt of N-phosphonomethyliminodiacetic acid, whereby the salt-forming cation is selected from the group alkali metals, alkaline earth metals, ammonium and organic ammonium, in the presence of oxygen in an aqueous-basic or aqueous solution and in the presence of a noble metal catalyst on a solid carrier, under excess pressure and at elevated temperature, the process being characterised by the fact that (a) the noble metal is selected from the group platinum, palladium or mixtures of palladium and platinum, and platinum or palladium individually or together in a mixture with ruthenium, rhodium, osmium or iridium, and (b) it is additionally admixed with at least one doping metal from the group bismuth, lead, tin, tellurium, germanium, vanadium, chromium, iron, manganese, molybdenum and tungsten. The known herbicide N-phosphonomethylglycine may be prepared from N-phosphonomethyliminodiacetic acid.</p> <p style="text-align: right;">Jerry R. Ebner, et al. Serial No. 09/408,323 Filed 09/29/99 Our File MTC 6610 Ref. No. 82</p>		

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	*KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Preparation of N-phosphonomethylglycine salts


The invention relates to a catalytic preparation process for N-phosphonomethylglycine salts by means of oxidative dealkylation of N-phosphonomethyliminodiacetic acid with oxygen in an aqueous-basic solution and in the presence of a noble metal catalyst on a solid carrier, the process being characterised by the fact that (a) the noble metal is selected from the group platinum, palladium or mixtures of palladium and platinum, and platinum or palladium individually or together in a mixture with ruthenium, rhodium, osmium or iridium, and (b) it is additionally mixed with at least one metal from the group bismuth, lead, tin, tellurium, germanium, vanadium, chromium, iron, manganese, molybdenum and tungsten.

N-phosphonomethylglycine of formula I



is a known herbicide or a known plant growth regulator, see US-A-4 147 719.

From US-A-4 147 719 a catalytic preparation process for N-phosphonomethylglycine salts is also known, in which a salt of N-phosphonomethyliminodiacetic acid of formula II



is reacted with oxygen in an aqueous-basic solution and in the presence of a platinum catalyst on activated carbon as a solid carrier. Disadvantages of this process are: (a) the use of a specific carrier that has to be activated, which increases the costs of the catalyst, (b) the restriction of the noble metal to platinum, and (c) the dissolving of substantial amounts of platinum in the product, which is linked with deactivation of the catalyst and thus limited reusability and platinum losses during reworking. In all, these disadvantages lead to an uneconomical process that cannot therefore be used in industry.

It has now surprisingly been found that (a) both platinum and palladium can be used either individually or mixed together and additionally in admixture with further noble metals, (b) non-activated solid carriers can be used, (c) no losses in catalyst activity or catalyst selectivity are observed, and (d) the losses in noble metals are substantially reduced, if at least one further metal is added from the group bismuth, lead, tin, tellurium, germanium, vanadium, chromium, iron, manganese, molybdenum and tungsten. As a rule, higher yields are achieved, and the catalysts may be reused several times.

The subject matter of the present invention is thus a process for the preparation of N-phosphonomethylglycine salts by reacting a salt of N-phosphonomethyliminodiacetic acid, whereby the salt-forming cation is selected from the group alkali metals, alkaline earth metals, ammonium and organic ammonium, in the presence of oxygen in an aqueous-basic or aqueous solution and in the presence of a noble metal catalyst on a solid carrier, under excess pressure and at elevated temperature, the process being characterised by the fact that (a) the noble metal is selected from the group platinum, palladium or mixtures of palladium and platinum, and platinum or palladium individually or together in a mixture with ruthenium, rhodium, osmium or iridium, and (b) it is additionally admixed with at least one doping metal from the group bismuth, lead, tin, tellurium, germanium, vanadium, chromium, iron, manganese, molybdenum and tungsten.

Preferred alkali metals are Li, Na, K, Rb and Cs, most preferably Na and K. Preferred alkaline earth metals are Mg, Ca, Ba and Sr. Organic ammonium is understood to be, in the context of the invention, salt-forming cations of primary, secondary, tertiary and quaternary ammonium. The N-atom may be substituted by aromatic radicals having 6 to 14 carbon atoms, aromatic-aliphatic radicals having 7 to 15 carbon atoms, or preferably linear or branched aliphatic radicals having 1 to 12 carbon atoms, or preferably cyclic aliphatic radicals having 3 to 12 ring carbon atoms. Especially preferred is organic ammonium with 1 to 4 identical or different substituents from the group linear or branched alkyl with 1 to 8, more preferably 1 to 6, and most preferably 1 to 4, carbon atoms, and cycloalkyl with having 5 or 6 ring carbon atoms. Primary C₁-C₆-alkylammonium is preferred in particular. Salt-forming cations are most preferably derived from metals selected from the group Na, K, Mg, Ca and Ba, as well as ammonium (NH₄⁺) and C₁-C₆-alkylammonium.

In order to form the salt, the corresponding alkali metal hydroxides, alkaline earth metal hydroxides, ammonium hydroxides or organic ammonium hydroxide are added to the aqueous solution of N-phosphonomethyliminodiacetic acid. The molar ratio of N-phosphonomethyliminodiacetic acid to hydroxide (or to salt-forming cation) may be from 1:1 to 1:2, preferably 1:1.5 and most preferably 1:1.3. The reaction mixture may accordingly be basic, neutral or acidic, depending on the molar ratio.

Oxygen may be used in pure form, or mixed with inert gases, for example nitrogen or noble gases. The oxygen is generally under pressure, so that the reaction is carried out under excess pressure, for example 1000 hPa to 10,000 hPa, preferably 1000 to 5000 hPa.

The reaction temperature may be for example 30 to 200° C, preferably 60 to 150° C, most preferably 80 to 130° C.

The solid carriers may be for example finely divided activated carbon, finely divided SiO₂ gel, SiO₂ mixed gel with Al, Ti and/or Zr oxides, finely divided silicate minerals (diatomaceous earth, alumina, zeolites, bentonites, montmorillonites) and finely divided aluminium oxide. The preferred carrier is activated carbon.

The catalyst metal is preferably present in a quantity of 1 to 10, more preferably 2 to 5 % by weight, based on the carrier. The metals used for doping may be present in an amount of 0.5 to 10, more preferably 1 to 5 % by weight. The weight ratio of noble metal or noble metals to doping metal is preferably 1:20 to 1:1, most preferably 1:10 to 1:1.

Preferred noble metals are Pd and especially Pt. Preferred doping metals are Bi, Pb and Sn, most preferably Bi and Pb. These preferred noble metals and doping metals are used in particular on activated carbon as the carrier. The catalysts that have proved to be particularly effective are those which contain platinum on activated carbon and the doping metals bismuth or lead, whereby platinum is present in an amount of 3 to 6 % by weight and the doping metal in an amount of 1 to 6 % by weight.

The catalyst consisting of carrier / noble metal / doping metal is preferably employed in amounts of 4 to 20, more preferably 4 to 15, most preferably 5 to 10 % by weight, based on the amount of N-phosphonomethyliminodiacetic acid.

The process may be carried out for example, whereby N-phosphonomethyliminodiacetic acid, the catalyst and the base are added to water in a pressurised container, and the mixture is heated to the reaction temperature. Then, the air is expelled by applying oxygen pressure, and the reaction mixture is stirred or shaken. The pressure is increased by about 500 to 1000 hPa through the formation of CO₂. Afterwards, the gas phase is drawn off, and oxygen pressure is again applied. When no further pressure increase is observed, the reaction is complete. The desired N-phosphonomethylglycine salt is isolated for example by evaporating off the water and the formaldehyde that has formed. The acid may be produced from the salt in conventional manner, by treatment with an acid.

In the process according to the invention, the yield is considerably higher than 90% and the N-phosphonomethylglycine (PMG) content in the isolated product is at least 70%. The noble metal content in the isolated product is less than 10 ppm, whereas in processes without the addition of a doping metal, the content is considerably higher than 200 ppm. In the isolated product, at first amounts of more than 500 ppm of doping metal are established. When the catalyst is reused, this amount is reduced substantially, depending on the number of reuses, to below 10 ppm. The catalyst may be reused up to ten times or more, with only a slight loss in activity being observed.

The catalysts are known, partly commercial, or may be produced by analogous processes, for example separating from corresponding metal salt solutions by reduction on the carrier.

The following examples illustrate the invention more fully.

Example 1:

a) Usage of a Pt/Bi catalyst on activated carbon

100 ml of water, 12 g (0.049 mols) of N-phosphonomethyliminodiacetic acid (NPDS) and 3.15 g (0.053 mols) of isopropylamine are placed in a pressurised container. Afterwards, 1.0 g of catalyst is added (3% each of Pt and Bi on activated carbon; Degussa CF 196 XR/W B93/11) and heated to 90-105°C. Then, oxygen pressure is applied to a pressure of

2000 hPa, the pressure is released and then reapplied with oxygen at 2000 hPa. The container is shaken during the reaction. After increasing the pressure to 2500 hPa, the gas phase is drawn off, 2 bars oxygen pressure are again applied, and shaking continues. The reaction is complete after 2 hours. After cooling, the catalyst is filtered off and the reaction mixture is evaporated to dryness. The yield is 10.7 g (96%) NPDS-isopropylamine salt; the content of N-phosphonomethylglycine (PMG) according to HPLC analysis is 84% and the content of Pt is only 4 ppm.

a) Usage of a Pt catalyst on activated carbon

Example a) is repeated and 1.0 g of catalyst (5% Pt on activated carbon) is used. The yield is 9.0 g (72%) NPDS-isopropylamine salt; the content of PMG according to HPLC analysis is 79% and the content of Pt is 210 ppm.

Example 2: Usage of a Pt/Pb catalyst.

Example 1a) is repeated and 1.0 g of catalyst (5% Pt and 1% Pb on activated carbon; Degussa CF 101 R/W) is used. The yield is 10.0 g (96%) NPDS-isopropylamine salt; the content of PMG according to HPLC analysis is 85% and the content of Pt is 5 ppm.

Examples 3-6 Reuse of the catalyst

Example 1a) is repeated and the catalyst used is 3% each of Pt and Bi on activated carbon; Degussa CF 196 XR/W B93/11 (examples 3 and 4). The separated catalyst is washed three times with deionised water and used again without further treatment (examples 5 and 6). Further details may be found in Table 1.

Example No.	catalyst (g)	yield (g) (% of theory)	PMG content % according to HPLC	ppm Pt in PMG	ppm Bi in PMG
3	1 (A)	11.0 (98)	78	4	840
4	0.08 (B)	10.6 (95)	85	6	740
5	A	10.4 (93)	85	6	230
6	B	10.8 (96)	80	5	33

What is claimed is:

1. Process for the preparation of N-phosphonomethylglycine salts by reacting a salt of N-phosphonomethyliminodiacetic acid, whereby the salt-forming cation is selected from the group alkali metals, alkaline earth metals, ammonium and organic ammonium, in the presence of oxygen in an aqueous-basic or aqueous solution and in the presence of a noble metal catalyst on a solid carrier, under excess pressure and at elevated temperature, the process being characterised by the fact that (a) the noble metal is selected from the group platinum, palladium or mixtures of palladium and platinum, and platinum or palladium individually or together in a mixture with ruthenium, rhodium, osmium or iridium, and (b) it is additionally admixed with at least one doping metal from the group bismuth, lead, tin, tellurium, germanium, vanadium, chromium, iron, manganese, molybdenum and tungsten.
2. Process according to claim 1, wherein the salt-forming cations are derived from metals selected from the group Na, K, Mg, Ca and Ba, as well as ammonium (NH_4^+) and $\text{C}_1\text{-C}_6$ -alkylammonium
3. Process according to claim 2, wherein the alkylammonium is isopropylamine.
4. Process according to claim 1, wherein the molar ratio of N-phosphonomethyliminodiacetic acid to salt-forming cation is 1:1 to 1:2.
5. Process according to claim 1, wherein the excess pressure is 1 to 10 bar.
6. Process according to claim 1, wherein the reaction temperature is 30 to 200°C.
7. Process according to claim 1, wherein the catalyst metal is present in a quantity of 1 to 10% by weight, based on the carrier, and the metals used for doping are present in an amount of 0.5 to 10% by weight.
8. Process according to claim 1, wherein the weight ratio of noble metal or noble metals to doping metal is 1:20 to 1:1.

9. Process according to claims 1 to 8, wherein the noble metal is Pt and the doping metals are Bi and Pb, and the carrier is activated carbon.

10. Process according to claim 9, wherein the catalyst is platinum on activated carbon and contains the doping metals bismuth or lead, whereby platinum is present in an amount of 3 to 6 % by weight and the doping metal in an amount of 1 to 6 % by weight, based on the carrier.

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07F9/38

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4 147 719 A (JOHN E. FRANZ) 3 April 1979 (1979-04-03) cited in the application the whole document ---	1-10
Y	US 3 950 402 A (JOHN E. FRANZ) 13 April 1976 (1976-04-13) the whole document ---	1-10
Y	EP 0 472 693 A (ERCROS S.A.) 4 March 1992 (1992-03-04) the whole document ---	1-10
Y	EP 0 098 034 A (GESHURI LABORATORIES LTD.) 11 January 1984 (1984-01-11) the whole document ---	1-10
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

8 November 1999

Date of mailing of the international search report

15/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Beslier, L

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	<p>WO 99 43430 A (MONSANTO CO.) 2 September 1999 (1999-09-02) page 22, line 17 -page 26, line 14; claims 1,4,7-9,281-379</p> <p>-----</p>	1-10

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4147719 A	03-04-1979	AR 220326 A	31-10-1980
		AT 364853 B	25-11-1981
		AT 906877 A	15-04-1981
		AU 515070 B	12-03-1981
		AU 3172577 A	28-06-1979
		BE 861996 A	19-06-1978
		BG 42358 A	14-11-1987
		BR 7708434 A	08-08-1978
		CA 1090819 A	02-12-1980
		CH 634580 A	15-02-1983
		CS 194697 B	31-12-1979
		DD 133952 A	31-01-1979
		DE 2756577 A	22-06-1978
		DK 565977 A, B,	21-06-1978
		EG 13039 A	31-07-1980
		ES 465139 A	01-10-1978
		FI 773835 A, B,	21-06-1978
		FR 2374328 A	13-07-1978
		GB 1575769 A	24-09-1980
		IE 46121 B	23-02-1983
		IL 53634 A	31-03-1981
		IT 1089971 B	18-06-1985
		JP 1141126 C	24-03-1983
		JP 53077018 A	08-07-1978
		JP 57029038 B	19-06-1982
		MX 4703 E	04-08-1982
		MY 12082 A	31-12-1982
		NL 7713959 A	22-06-1978
		PH 14143 A	05-03-1981
		SE 436282 B	26-11-1984
		SE 7714408 A	21-06-1978
		YU 301377 A	30-06-1982
		ZA 7707533 A	25-10-1978
US 3950402 A	13-04-1976	US 3954848 A	04-05-1976
		AU 5627773 A	05-12-1974
		BE 800238 A	30-11-1973
		BG 21212 A	20-03-1976
		CA 992093 A	29-06-1976
		CH 574968 A	30-04-1976
		CS 177133 B	29-07-1977
		DD 104303 A	05-03-1974
		DE 2327680 A	13-12-1973
		DE 2366060 A	06-10-1977
		DK 138951 B	20-11-1978
		EG 11140 A	15-08-1977
		ES 415335 A	01-05-1976
		ES 440610 A	01-03-1977
		FR 2186477 A	11-01-1974
		FR 2193831 A	22-02-1974
		GB 1432043 A	14-04-1976
		IE 37730 B	28-09-1977
		IL 42393 A	31-12-1975
		IT 988869 B	30-04-1975
		JP 49048620 A	11-05-1974
		JP 51011093 B	08-04-1976
		NL 7307449 A, B,	04-12-1973
		RO 62901 A	15-08-1977

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 3950402 A ¹		RO 66031 A	15-07-1980
		SE 404019 B	18-09-1978
		SE 407579 B	02-04-1979
		SE 7600415 A	16-01-1976
		YU 142773 A	18-06-1982
		YU 262880 A	30-06-1983
		AT 327946 B	25-02-1976
		AT 477373 A	15-05-1975
		ZA 7303705 A	24-04-1974

EP 472693 A	04-03-1992	ES 2021229 A	16-10-1991
		DE 69114610 D	21-12-1995
		DE 69114610 T	02-05-1996
		GR 3018731 T	30-04-1996
		AU 637896 B	10-06-1993
		AU 7497391 A	10-10-1991
		DK 472693 T	02-01-1996
		WO 9113893 A	19-09-1991
		US 5179228 A	12-01-1993

EP 98034 A	11-01-1984	IL 66137 A	31-03-1986
		AU 1350283 A	05-01-1984
		BR 8303513 A	07-02-1984
		DK 157683 A	26-12-1983
		FI 831204 A	26-12-1983
		GR 81421 A	11-12-1984
		JP 59007196 A	14-01-1984
		PH 18543 A	09-08-1985
		PT 76635 A, B	01-06-1983
		RO 87143 A	30-11-1985
		US 4486356 A	04-12-1984
		ZA 8302562 A	27-06-1984
		ES 523125 A	01-10-1984

WO 9943430 A	02-09-1999	NONE	
